

***Amendments to the Specification***

Please replace the paragraph beginning on page 15, line 10 with the following paragraph:

A complex of an HLA molecule and a peptidic antigen acts as the ligand recognized by HLA-restricted T cells (Buus, S *et al.*, *Cell* 47:1071, 1986; Babbitt, B.P. *et al.*, *Nature* 317:359, 1985; Townsend, A. and Bodmer, H. *Annu. Rev. Immunol.* 7:601, 1989; Germain, R.N., *Annu. Rev. Immunol.* 11:403, 1993). Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues that correspond to motifs required for specific binding to HLA antigen molecules have been identified and are described herein and are set forth in Tables I, II, and III (*see also, e.g.,* Southwood, *et al.*, *J. Immunol.* 160:3363, 1998; Rammensee, *et al.*, *Immunogenetics* 41:178, 1995; Rammensee *et al.*, SYFPEITHI, ~~access via web at: [http://](http://134.2.96.221/scripts.hlaserver.dll/home.htm)~~ which can be found on the World Wide Web at the URL 134.2.96.221/scripts.hlaserver.dll/home.htm; Sette, A. and Sidney, *J. Curr. Opin. Immunol.* 10:478, 1998; Engelhard, V.H., *Curr. Opin. Immunol.* 6:13, 1994; Sette, A. and Grey, H.M., *Curr. Opin. Immunol.* 4:79, 1992; Sinigaglia, F. and Hammer, *J. Curr. Biol.* 6:52, 1994; Ruppert, *et al.*, *Cell* 74:929-937, 1993; Kondo *et al.*, *J. Immunol.* 155:4307-4312, 1995; Sidney *et al.*, *J. Immunol.* 157:3480-3490, 1996; Sidney *et al.*, *Human Immunol.* 45:79-93, 1996; Sette, A. and Sidney, *J. Immunogenetics*, in press, 1999).